Sample results. Actual results may vary.

PATIENT INFORMATION

Clinical Info:

SPECIMEN INFORMATION

SPECIMEN:
REQUISITION:
LAB REF NO:

DOB:
AGE:
GENDER:
FASTING:

COLLECTED:
RECEIVED:
REPORTED:

REPORT STATUS: FINAL

ORDERING PHYSICIAN



IU/mL

IU/mL

Order Today

www.accesalabs.com/titers

Test Name Result Flag Reference Range Lab

BORDETELLA PERTUSSIS IGG ABS, MAID

PT IGG <1 FHA IGG 13

> REFERENCE RANGES: PT IgG <45 IU/mL FHA IgG <90 IU/mL

Levels of antibodies recognizing pertussis toxin (PT) and filamentous hemagglutinin (FHA) are typically increased following vaccination or natural exposure to Bordetella pertussis. This

1 of 2

ΧE

XE

Sample results. Actual results may vary.

assay is not appropriate for assessing immunity to pertussis because the specific antibodies and antibody levels that correlate with protection have not been well defined. The indicated reference range values reflect the 95th percentile of antibody levels from blood donors; thus, antibody levels above the reference range are highly suggestive of recent infection or vaccination. Increased levels of FHA antibodies alone may represent cross-reactive antibodies induced by infection with other Bordetella species, Mycoplasma pneumoniae, Chlamydophila pneumoniae, or nonencapsulated Haemophilus influenzae.

