

Sample results. Actual results may vary.

PATIENT INFORMATION

REPORT STATUS: FINAL

ORDERING PHYSICIAN

CLIENT INFORMATION



Order Today

www.accesalabs.com/arthritis

SPECIMEN INFORMATION

SPECIMEN:

DOB:

REQUISITION:

AGE:

LAB REF NO:

GENDER:

FASTING:

COLLECTED:

Clinical Info:

RECEIVED:

REPORTED:

Test Name	Result	Flag	Reference Range	Lab
ANCA VASCULITIDES				
PROTEINASE-3 ANTIBODY	<1.0		<1.0 AI	01

Value Interpretation
 <1.0 AI: No Antibody Detected
 >or=1.0 AI: Antibody Detected

Autoantibodies to proteinase-3 (PR-3) are accepted as characteristic for granulomatosis with polyangiitis (Wegener's), and are detectable in 95% of the histologically proven cases. The cytoplasmic IFA pattern, (c-ANCA), is based largely on autoantibody to PR-3 which serves as the primary antigen.

These autoantibodies are present in active disease state.

MYELOPEROXIDASE ANTIBODY	<1.0		<1.0 AI	01
--------------------------	------	--	---------	----

Value Interpretation
 <1.0 AI: No Antibody Detected
 >or=1.0 AI: Antibody Detected

Autoantibodies to myeloperoxidase (MPO) are commonly associated with the following small-vessel vasculitides: microscopic polyangiitis, polyarteritis nodosa, Churg-Strauss syndrome, necrotizing and crescentic glomerulonephritis and occasionally Wegener's granulomatosis. The perinuclear IFA pattern, (p-ANCA) is based largely on autoantibody to myeloperoxidase which serves as the primary antigen.

These autoantibodies are present in active disease state.

