

**SAMPLE RESULTS. ACTUAL RESULTS MAY VARY.**

PATIENT INFORMATION

REPORT STATUS: FINAL

ORDERING PHYSICIAN

CLIENT INFORMATION



ORDER TODAY

SPECIMEN INFORMATION

SPECIMEN:

REQUISITION:

LAB REF NO:

DOB:

AGE:

GENDER:

FASTING:

Clinical Info:

COLLECTED:

RECEIVED:

REPORTED:

Test Name	Result	Flag	Reference Range	Lab
FASTING: UNKNOWN				
<b>GLUCOSE</b>				
GLUCOSE	133	HIGH	65-99 mg/dL	CB
Fasting reference interval				
<b>HEMOGLOBIN A1c</b>				
HEMOGLOBIN A1c	6.8	HIGH	<5.7 % of total Hgb	CB
<p>According to ADA guidelines, hemoglobin A1c &lt;7.0% represents optimal control in non-pregnant diabetic patients. Different metrics may apply to specific patient populations. Standards of Medical Care in Diabetes-2013. Diabetes Care. 2013;36:s11-s66</p> <p>For the purpose of screening for the presence of diabetes</p> <p>&lt;5.7% Consistent with the absence of diabetes</p> <p>5.7-6.4% Consistent with increased risk for diabetes (prediabetes)</p> <p>&gt;or=6.5% Consistent with diabetes</p> <p>This assay result is consistent with diabetes mellitus.</p> <p>Currently, no consensus exists for use of hemoglobin A1c for diagnosis of diabetes for children.</p>				

**Performing Laboratory Information:**