

Sample results. Actual results may vary.

PATIENT INFORMATION

REPORT STATUS: FINAL

ORDERING PHYSICIAN

CLIENT INFORMATION



Order Today

www.accessalabs.com/thyroid

SPECIMEN INFORMATION

SPECIMEN:
REQUISITION:
LAB REF NO:

DOB:
AGE:
GENDER:
FASTING:

Clinical Info:

COLLECTED:
RECEIVED:
REPORTED:

| Test Name | Result | Flag | Reference Range | Lab |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|------|-----------------|-----|
| TSI (THYROID STIMULATING IMMUNOGLOBULIN) | | | | |
| TSI | <89 | | <140 % baseline | 01 |
| <p>Thyroid stimulating immunoglobulins (TSI) can engage the TSH receptors resulting in hyperthyroidism in Graves' disease patients. TSI levels can be useful in monitoring the clinical outcome of Graves' disease as well as assessing the potential for hyperthyroidism from maternal-fetal transfer. TSI results greater than or equal to (>=) 140% of the Reference Control are considered positive.</p> <p>NOTE: A serum TSH level greater than 350 micro-International Units/mL can interfere with the TSI bioassay and potentially give false positive results.</p> <p>Patients who are pregnant and are suspected of having hyperthyroidism should have both a TSI and human Chorionic Gonadotropin (hCG) tests measured. A serum hCG level greater than 40,625 mIU/mL can interfere with the TSI bioassay and may give false negative results. In these patients it is recommended that a second TSI is obtained when the hCG concentration falls below 40,625 mIU/mL (usually after approximately 20-weeks gestation).</p> | | | | |
| THYROID PEROXIDASE ANTIBODIES | | | | |
| THYROID PEROXIDASE ANTIBODIES | 1 | | <9 IU/mL | 02 |
| T3, TOTAL | | | | |
| T3, TOTAL | 59 | LOW | 76-181 ng/dL | 02 |
| T4, FREE | | | | |
| T4, FREE | 1.4 | | 0.8-1.8 ng/dL | 02 |
| T4 (THYROXINE), TOTAL | | | | |
| T4 (THYROXINE), TOTAL | 10.7 | | 4.5-12.0 mcg/dL | 02 |
| FREE T4 INDEX (T7) | DNR | | 1.4-3.8 | 02 |
| TSH | | | | |
| TSH | 2.80 | | 0.40-4.50 mIU/L | 02 |
| T3, FREE | | | | |
| T3, FREE | 1.7 | LOW | 2.3-4.2 pg/mL | 02 |

Performing Laboratory Information: